

# Publications

## Whole-transcript analysis

The following publications and articles demonstrate the advantages of GeneChip® Gene 1.0 ST and Exon 1.0 ST Arrays for whole-transcript analysis. The uses described in these reference articles have not been approved or cleared by the FDA.

### 2010

Aghajanova L., et al. The protein kinase A pathway-regulated transcriptome of endometrial stromal fibroblasts reveals compromised differentiation and persistent proliferative potential in endometriosis. *Endocrinology* **151**(3):1341-55 (2010).

Bradford J. R., et al. A comparison of massively parallel nucleotide sequencing with oligonucleotide microarrays for global transcription profiling. *BMC Genomics* **11**(1):282 (2010).

Chen H., et al. Differential expression and alternative splicing of genes in lumbar spinal cord of an amyotrophic lateral sclerosis mouse model. *Brain Research* **1340**:52-69 Epub ahead of print (2010).

Ciobanu D. C., et al. Detection, validation, and downstream analysis of allelic variation in gene expression. *Genetics* **184**(1):119-28 (2010).

de la Grange P., Gratadou L., Delord M., Dutertre M., Auboeuf D. Splicing factor and exon profiling across human tissues. *Nucleic Acids Research* **38**(9):2825-38 (2010).

Dutertre M., et al. Exon-based clustering of murine breast tumor transcriptomes reveals alternative exons whose expression is associated with metastasis. *Cancer Research* **70**(3):896-905 (2010).

Emig D., et al. AltAnalyze and DomainGraph: analyzing and visualizing exon expression data. *Nucleic Acids Research* **38**(Suppl):W755-62 Epub (2010).

Gamazon E. R., Zhang W., Dolan M. E., Cox N. J. Comprehensive survey of SNPs in the Affymetrix Exon Array using the 1000 Genomes dataset. *PLoS One* **5**(2):e9366 (2010).

Jansen B. J. H., et al. Functional differences between mesenchymal stem cell populations are reflected by their transcriptome. *Stem Cells and Development* **19**(4):481-90 (2010).

Kong S. W., et al. Heart failure-associated changes in RNA splicing of sarcomere genes. *Circulation: Cardiovascular Genetics* **3**(2):138-46 (2010).

Lin C. L., Evans V., Shen S., Xing Y., Richter J. D. The nuclear experience of CPEB: Implications for RNA processing and translational control. *RNA* **16**(2):338-48 (2010).

Lin H. Q., et al. Differential gene expression profiling on the muscle of acetylcholinesterase knockout mice: A preliminary analysis. *Chemico-Biological Interactions* **187**(1-3):120-23 (2010).

Lin L., et al. Evolution of alternative splicing in primate brain transcriptomes. *Human Molecular Genetics* **19**(15):2958-73 (2010).

Mojica W., Hawthorn L. Normal colon epithelium: a dataset for the analysis of gene expression and alternative splicing events in colon disease. *BMC Genomics* **11**:5 (2010).

Nurtdinov R. N., Vasiliev M. O., Ershova A. S., Lossev I. S., Karyagina A. S. PLANdbAfy: probe-level annotation database for Affymetrix expression microarrays. *Nucleic Acids Research* **38**(Database issue): D726-30 (2010).

- Ortis F., et al. Cytokines interleukin-1 $\beta$  and tumor necrosis factor- $\alpha$  regulate different transcriptional and alternative splicing networks in primary  $\beta$ -cells. *Diabetes* **59**(2):358-74 (2010).
- Qu Y., He F., Chen Y. Different effects of the probe summarization algorithms PLIER and RMA on high-level analysis of Affymetrix Exon Arrays. *BMC Bioinformatics* **11**:211 (2010).
- Rasche A., Herwig R. ARH: predicting splice variants from genome-wide data with modified entropy. *Bioinformatics* **26**(1):84-90 (2010).
- Ren H., et al. Characterization of peroxisome proliferator-activated receptor alpha-independent effects of PPARalpha activators in the rodent liver: di-(2-ethylhexyl) phthalate also activates the constitutive-activated receptor. *Toxicological Sciences* **113**(1):45-59 (2010).
- Richard, H., et al. Prediction of alternative isoforms from exon expression levels in RNA-Seq experiments. *Nucleic Acids Research* **38**(10):e112 (2010).
- Risueño A., Fontanillo C., Dinger M. E., De Las Rivas J. GATEExplorer: genomic and transcriptomic explorer; mapping expression probes to gene loci, transcripts, exons and ncRNAs. *BMC Bioinformatics* **11**:221 (2010).
- Rollins B., Martin M. V., Morgan L., Vawter M. P. Analysis of whole genome biomarker expression in blood and brain. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics* **153B**(4):919-36 (2010).
- Salomonis N., et al. Alternative splicing regulates mouse embryonic stem cell pluripotency and differentiation. *Proc Natl Acad Sci USA* **107**(23):10514-19 (2010).
- Seitz G., et al. Inhibition of glutathione-S-transferase as a treatment strategy for multidrug resistance in childhood rhabdomyosarcoma. *International Journal of Oncology* **36**(2):491-500 (2010).
- Shehadeh L. A., et al. SRRM2, a potential blood biomarker revealing high alternative splicing in Parkinson's disease. *PLoS One* **5**(2):e9104 (2010).
- Shelton R. C., et al. Altered expression of genes involved in inflammation and apoptosis in frontal cortex in major depression. *Molecular Psychiatry* Epub ahead of print (2010).
- Shen S., Warzecha C. C., Carstens R. P., Xing Y. MADS+: discovery of differential splicing events from Affymetrix exon junction array data. *Bioinformatics* **26**(2):268-9 (2010).
- Subbaram S., Kuentzel M., Frank D., Dipersio C. M., Chittur S. V. Determination of alternate splicing events using the Affymetrix Exon 1.0 ST Arrays. *Methods in Molecular Biology* **632**:63-72 (2010).
- Turro E., et al. MMBGX: a method for estimating expression at the isoform level and detecting differential splicing using whole-transcript Affymetrix arrays. *Nucleic Acids Research* **38**(1):e4 (2010).
- Wasim M., et al. PLZF/ZBTB16, a glucocorticoid response gene in acute lymphoblastic leukemia, interferes with glucocorticoid-induced apoptosis. *The Journal of Steroid Biochemistry and Molecular Biology* **120**(4-5):218-27 (2010).
- Whistler T., Chiang C. F., Lin J. M., Lonergan W., Reeves W. C. The comparison of different pre- and post-analysis filters for determination of exon-level alternative splicing events using Affymetrix arrays. *Journal of Biomolecular Techniques* **21**(1):44-53 (2010).
- Younis I., et al. Rapid-response splicing reporter screens identify differential regulators of constitutive and alternative splicing. *Molecular and Cellular Biology* **30**(7):1718-28 (2010).
- Zhang Z., Gasser D. L., Rappaport E. F., Falk M. J. Cross-platform expression microarray performance in a mouse model of mitochondrial disease therapy. *Molecular Genetics and Metabolism* **99**(3):309-18 (2010).

## 2009

Cheung H. C., et al. Splicing factors PTBP1 and PTBP2 promote proliferation and migration of glioma cell lines. *Brain* **132**(8):2277-88 (2009).

Dolnik A., et al. Genome-wide analysis of alternative splicing points to novel leukemia relevant genes in acute Myeloid Leukemia. *Blood* (ASH Annual Meeting Abstracts) **114**:2391 (2009).

Gaidatzis D., Jacobait K., Oakeley E. J., Stadler M. B. Overestimation of alternative splicing caused by variable probe characteristics in exon arrays. *Nucleic Acids Research* **37**(16):e107 (2009).

Gellert P., Uchida S., Braun T. Exon Array Analyzer: a web interface for Affymetrix Exon Array analysis. *Bioinformatics* **25**(24):3323-4 (2009).

Gillett A., et al. Alternative splicing and transcriptome profiling of experimental autoimmune encephalomyelitis using genome-wide exon arrays. *PLoS One* **4**(11):e7773 (2009).

Ha K. C., Coulombe-Huntington J., Majewski J. Comparison of Affymetrix Gene ST Array with the Exon Array shows potential application for detection of transcript isoform variation. *BMC Genomics* **10**:519 (2009).

Hang X., et al. Transcription and splicing regulation in human umbilical vein endothelial cells under hypoxic stress conditions by exon array. *BMC Genomics* **10**:126 (2009).

Howard B. E., Sick B., Heber S. Unsupervised assessment of microarray data quality using a Gaussian mixture model. *BMC Bioinformatics* **10**:191 (2009).

Hu J., He X., Cote G. J., Krahe R. J. Singular value decomposition-based alternative splicing detection. *Journal of the American Statistical Association* **104**(487):944-53 (2009).

Huang G. J., et al. High resolution mapping of expression QTLs in heterogeneous stock mice in multiple tissues. *Genome Research* **19**(6):1133-40 (2009).

Jhavar S., et al. Integration of ERG gene mapping and gene-expression profiling identifies distinct categories of human prostate cancer. *BJU International* **103**(9):1256-69 (2009). Comment in: *BJU International* **103**(9):1293; author reply **103**(9):1293 (2009).

Juul S. E., et al. Microarray analysis of high-dose recombinant erythropoietin treatment of unilateral brain injury in neonatal mouse hippocampus. *Pediatric Research* **65**(5):485-92 (2009).

Laajala E., Aittokallio T., Lahesmaa R., Elo L. L. Probe-level estimation improves the detection of differential splicing in Affymetrix Exon Array studies. *Genome Biology* **10**(7):R77 (2009).

Lacaze P., et al. Combined genome-wide expression profiling and targeted RNA interference in primary mouse macrophages reveals perturbation of transcriptional networks associated with interferon signalling. *BMC Genomics* **10**(1):372 (2009).

Lin L., et al. Using high-density exon arrays to profile gene expression in closely related species. *Nucleic Acids Research* **37**(12):e90 (2009).

Linton K. M., et al. Methods comparison for high-resolution transcriptional analysis of archival material on Affymetrix Plus 2.0 and Exon 1.0 microarrays. *Biotechniques* **47**(1):587-96 (2009).

Liu L., et al. Age-associated changes in cardiac gene expression after preconditioning. *Anesthesiology* **111**(5):1052-64 (2009).

Miki L., et al. Alternative pre-mRNA splicing switches modulate gene expression in late erythropoiesis. *Blood* **113**(14):3363-70 (2009).



Morsczeck C., et al. Gene expression profiles of dental follicle cells before and after osteogenic differentiation in vitro. *Clinical Oral Investigations* **13**(4):383-91 (2009).

Pitroda S., et al. STAT1-dependent expression of energy metabolic pathways links tumour growth and radioresistance to the Warburg effect. *BMC Medicine* **7**:68 (2009).

Pohl A. A., et al. Affy exon tissues: exon levels in normal tissues in human, mouse and rat. *Bioinformatics* **25**(18):2442-3 (2009).

Robinson M. D., Speed T. P. Differential splicing using whole-transcript microarrays. *BMC Bioinformatics* **10**:156 (2009).

Shah S. H., Pallas J. A. Identifying differential exon splicing using linear models and correlation coefficients. *BMC Bioinformatics* **10**:26 (2009).

Wang X., et al. Genome-wide prediction of cis-acting RNA elements regulating tissue-specific pre-mRNA alternative splicing. *BMC Genomics* **10** Suppl 1:S4 (2009).

Warzecha C. C., Shen S., Xing Y., Carstens R. P. The epithelial splicing factors ESRP1 and ESRP2 positively and negatively regulate diverse types of alternative splicing events. *RNA Biology* **6**(5):546-62 (2009).

Weigelt K., Lichtinger M., Rehli M., Langmann T. Transcriptomic profiling identifies a PU.1 regulatory network in macrophages. *Biochemical and Biophysical Research Communications* **380**(2):308-12 (2009).

Zhang W., et al. Identification of common genetic variants that account for transcript isoform variation between human populations. *Human Genetics* **125**(1):81-93 (2009).