## **invitrogen** Human IL-8 Singleplex Bead Kit Technical Data Sheet

Catalog #: LHC0081 Lot #:\* 953118

\*Note: A letter at the end of the lot number signifies an additional packaging of this same lot.

### **Intended Use**

This kit is comprised of components for the measurement of human IL-8 in serum, plasma or tissue culture supernatant. The assay may be run alone or in combination with other Singleplex Bead Kits from Invitrogen. Buffer reagents needed to complete the reaction are sold separately under Catalog #LHB0001. These reagents are intended for use in the Luminex® 100<sup>TM</sup>/200<sup>TM</sup> and the FlexMAP 3D® System. This kit is configured for research use only and is not to be used in diagnostic procedures.

**TT TT** 0

### **Reagents Provided**

				Hu IL-8				
1.	Part #:	LM014	Description:	Antibody Bead Concentrate (10X)	Lot:	985071	Size:	0.25 mL-100 tests
Bead Region: Form: Storage:		81 0.25 mL 10X bead concentrate solution in storage buffer. Contains 0.05% sodium azide as a preservative. <b>Light-sensitive material.</b> Store at 2 to 8°C in the dark, until the expiration date indicated on the kit.						
2.	Part #:	BN014	Description:	Hu IL-8 Biotinylated Ab Conc. (10X)	Lot:	985069	Size:	1 mL-100 tests
Form: Storage:		1 mL of a 10X stock of Biotinylated Antibody Concentrate in Biotin Diluent. Contains 0.1% sodium azide as preservative. Concentration of antibody is matched to this lot of beads. Do not mix lots of Coated Beads and Detection Antibody. Store at 2 to 8°C until the expiration date indicated on the kit.						
3.	Part #:	SM311	Description:	Hu 16-Plex Standard	Lot:	925003	Size:	2 Vials
<ul> <li>Form: Lyophilized. The proteins in this standard have been calibrated against the masses of highly purified recomb proteins, with the respective Invitrogen ELISA kits, and NIBSC calibration standards (if available). Refer to the Manual included in the Buffer Reagent Kit for further information. Contains 0.1% sodium azide as a preservativ Store at 2 to 8°C. Use within 1 hour after reconstitution. Discard immediately after use.</li> </ul>			ole). Refer to the User					
Concentration of Reconstituted Standard**:								

IL-1β (8,200 pg/mL)	IL-10 (19,700 pg/mL)	IFN-α (13,800 pg/mL)	IL-6 (5,150 pg/mL)
IL-12 (13,400 pg/mL)	IL-13 (21,700 pg/mL)	IL-15 (26,600 pg/mL)	IL-17 (25,600 pg/mL)
GM-CSF (19,200 pg/mL)	IL-5 (5,250 pg/mL)	IFN-γ (16,200 pg/mL)	TNF-α (11,100 pg/mL)
IL-2 (18,400 pg/mL)	IL-7 (6,700 pg/mL)	IL-4 (49,000 pg/mL)	IL-8 (11,000 pg/mL)

## **\*\*Important note:** The concentrations of reconstituted standard are lot-specific. Please verify all concentration values entered in data analysis software.

One microgram of Invitrogen recombinant human IL-8 equals 1000 International Units of WHO reference preparation 89/520 (NIBSC, Hertfordshire, UK, EN6 3QG).

**Reconstitution:** Reconstitute with 1 mL Assay Diluent when measuring IL-8 in serum or plasma samples. For other sample types, such as tissue culture supernatants, reconstitute the standard in 1 mL of a solution consisting of 50% Assay Diluent + 50% of sample matrix. Allow standard to rehydrate for approximately 10 minutes before further dilution.

**Recommended Starting Concentration for Standard Curve:** Upon reconstitution, the starting concentration of standard is the value cited above. Make serial 1:3 dilutions in Assay Diluent (serum/plasma samples) or other appropriate matrix. Use 100 µL per assay. If establishing a Multiplex Assay, this same standard can be used to measure the other related cytokines cited above in a Multiplex Assay format. Refer to the User Manual included in the Buffer Reagent Kit for further information.

### For Research Use Only. CAUTION: Not for human or animal therapeutic or diagnostic use.

www.invitrogen.com

Invitrogen Corporation • 7335 Executive Way, Frederick, MD 21704 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

(Page 1 of 2) MAN0004444 Rev 1.00 Effective Date: 01 JUN 2011

#### LHC0081

Limited Use Label License: Research Use Only

The purchase of this product conveys to the purchaser the limited, non-transferable right to use the purchased amount of the product only to perform internal research for the sole benefit of the purchaser. No right to resell this product or any of its components is conveyed expressly, by implication, or by estoppel. This product is for internal research purposes only and is not for use in commercial applications of any kind, including, without limitation, quality control and commercial services such as reporting the results of purchaser's activities for a fee or other form of consideration. For information on obtaining additional rights, please contact outlicensing@lifetech.com or Out Licensing, Life Technologies, 5791 Van Allen Way, Carlsbad, California 92008.


# Human IL-8 Singleplex Bead Kit

### **Technical Data Sheet**

### **Performance Characteristics**

**Analytical Sensitivity:** The minimum detectable dose of Hu IL-8 is <3 pg/mL. This was determined by adding two standard deviations to the mean FI obtained when the zero standard was assayed 30 times.

### **Typical Standard Curve**



**Specificity:** Buffered solutions of a panel of substances at 10 or 20 ng/mL were assayed with the Invitrogen Human IL-8 Singleplex Bead Kit. The following substances were tested and all were found to have no cross-reactivity: human IL-1 $\beta$ , IL-2, IL-4, IL-5, IL-6, IL-10, GM-CSF, IFN- $\gamma$ , MIP-2, TNF- $\alpha$ ; mouse IL-1 $\beta$ , IL-2, IL-4, IL-5, IL-6, IL-10, GM-CSF, IFN- $\gamma$ , MIP-2, TNF- $\alpha$ . Swine IL-8 showed 1% cross-reactivity.

### **Precision:**

	Intra-assay	Inter-assay
	(n=16)	(n=32)
Mean (pg/mL)	1020	987
SD	73	97
%CV	7.2	9.8

**Linearity:** Human serum and tissue culture medium containing 10% fetal calf serum were spiked with human IL-8 and serially diluted in Assay Diluent and a solution consisting of 50% Assay Diluent and 50% tissue culture medium containing 10% fetal calf serum, respectively, over the range of the assay. Linear regression analysis of samples versus the expected concentration yielded correlation coefficients of 0.98 and 0.99 for serum and tissue culture, respectively.

<b>Recovery:</b>
------------------

Sample type	Results
Serum (Human)	$\checkmark$
EDTA plasma (Human)	+
Citrate plasma (Human)	$\checkmark$
Heparin plasma (Human)	$\checkmark$
Tissue culture medium with 10% fetal calf serum*	$\checkmark$

Explanation of symbols				
Symbol	Description	Symbol	Description	
REF	Catalogue Number	LOT	Batch code	
RUO	Research Use Only	IVD	In vitro diagnostic medical device	
X	Use by	ł	Temperature limitation	
***	Manufacturer	EC REP	European Community authorised representative	
[-]	Without, does not contain	[+]	With, contains	
Î Î	Protect from light	$\triangle$	Consult accompanying documents	
i	Directs the user to consult instructions for use (IFU), accompanying the product.		or use (IFU), accompanying the product.	

**Notes:** 70-130% recovery ( $\checkmark$ ), 50-69% recovery (-), 131 -150% recovery (+) and <50% or >150% recovery (NR - Not recommended) \*Analysis performed during product development and with first lots produced.

### **Correlation to ELISA:**

A correlation coefficient of 0.97 was calculated when values for natural samples obtained with the Human IL-8 Singleplex Bead Kit were compared to the Invitrogen ELISA for Human IL-8 (cat.# KAC1301, KAC1302). Human IL-8 Singleplex Bead Kit (pg/mL) x 1.03 = Hu IL-8 ELISA (pg/mL).

### For Research Use Only. CAUTION: Not for human or animal therapeutic or diagnostic use.

www.invitrogen.com

Invitrogen Corporation • 7335 Executive Way, Frederick, MD 21704 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com

(Page 2 of 2) MAN0004444 Rev 1.00 Effective Date: 01 JUN 2011

### LHC0081

Limited Use Label License: Research Use Only

The purchase of this product conveys to the purchaser the limited, non-transferable right to use the purchased amount of the product only to perform internal research for the sole benefit of the purchaser. No right to resell this product or any of its components is conveyed expressly, by implication, or by estoppel. This product is for internal research purposes only and is not for use in commercial applications of any kind, including, without limitation, quality control and commercial services such as reporting the results of purchaser's activities for a fee or other form of consideration. For information on obtaining additional rights, please contact outlicensing@lifetech.com or Out Licensing, Life Technologies, 5791 Van Allen Way, Carlsbad, California 92008.