OXOID PRODUCT SPECIFICATION

PEPTONE WATERBO0208BTypical Formulagrams per litrePeptone10.0

Preparation

Sodium chloride

Suspend Peptone Water (15 grams / litre) in de-ionised water. Heat to dissolve. Dispense 3ml into final containers, bijou bottles. Sterilise at 121°C for 15 minutes. When cooled label each bottle and pack in units of 24 into labelled boxes.

5.0

Format

Twenty four bijou bottles with plastic screw cap closures in a box.

Labels

Label gives details of product name, product code, recommended storage temperature, lot number and expiry date.

Physical Characteristics

Physical Tests	
pН	7.5 ± 0.2
Colour	Straw
Clarity	Clear
Fill weight	3.0g + 0.3g

Packaging and presentation:

General appearance of bottle and label should be satisfactory. Label data should be correct.

Sterility Test

Macroscopic examination should show no evidence of microbial growth after incubation at 20 - 24°C and 30 - 34°C for 5 days.

Microbiological Tests Using Optimum Inoculum Dilution

Results after incubation at 35 - 39°C for 18 - 24 hours

Inoculum 10 -100 colony forming units.

Escherichia coli	ATCC [®] 25922	Turbid growth
Staphylococcus aureus	ATCC [®] 25923	Turbid growth

Storage conditions

Store away from light at between 2-25°C.